

# Monoclonal Anti-human IL-7R $\alpha$ /CD127

**Product reference: DDX0700**

## Description

The IL7-R consists of 2 chains, IL-7R $\alpha$  known as CD127 and common cytokine receptor  $\gamma$  chain known as CD132. A 75 to 80kDa human IL-7 receptor has been cloned that belongs to hematopoietic cytokine-receptor super family. R34-34, raised against human leukemic pre-B cells, recognized a molecule expressed on normal B cell precursors but not on mature B cells. This antibody specifically reverted IL-7 mediated growth inhibition of leukemic BCP (normal B cells precursors) and mature T cells. IL-7R $\alpha$  expression is dramatically influenced by cytokines and antigens. This IL-7R displays both high and low affinity for its ligand (IL-7). Inhibitory and proliferative effects of IL-7 can be mediated through the same receptor on various lineages. CD4<sup>+</sup> memory T cells express high level of IL-7R Subsets that express it generally require it, including progenitors of T and B cells, naïve and memory T cells.


(Pandrau-Garcia D et al, 1994, Blood, 83, 3613-9 ; Mazzucchelli R et al, Nat. Review Immunol., 2007,7, 144-54)

<b>Clone:</b>	<b>R34-34</b>
<b>Species:</b>	mouse
<b>Specificity:</b>	human IL7-R $\alpha$
<b>Immunogen:</b>	human leukemic pre-B cells
<b>Species cross- reactivity:</b>	dog
<b>Isotype:</b>	IgG1
<b>Formulation/size:</b>	<b>Purified:</b> 100 $\mu$ g in 200 $\mu$ l / 50 $\mu$ g in 100 $\mu$ l Tris-NaCl pH 8 <b>Coupled:</b> 100 $\mu$ g in 200 $\mu$ l / 50 $\mu$ g in 100 $\mu$ l PBS 50% glycerol


## Available formats:

Reference N°		Format	Application tested
50 $\mu$ g	100 $\mu$ g		
DDX0700P-50	DDX0700P-100	Purified	Surface Flow cytometry, IL7- binding inhibition, IP
DDX0700A488-50	DDX0700A488-100	Alexa-fluor® 488	Surface Flow cytometry
DDX0700A546-50	DDX0700A546-100	Alexa- fluor® 546	IF
DDX0700A647-50	DDX0700A647-100	Alexa- fluor® 647	Surface Flow cytometry


## Applications tested



Facs staining of R34-34-transfected Cos7 cells



Facs staining of PBL with R34-34



Immunoprecipitation of R34-34 antigen from <sup>125</sup>I-surface labelled Pre-Alp cells

## Usage recommendation:

- \*This monoclonal antibody may be used between 5-20  $\mu$ g/ml
- \*Optimal dilution should be determined by each laboratory for each application
- \*Coupled antibody: to maintain RT before using

## Aliquot storage conditions

- 20°C. KEEP CONTENTS STERILE: no preservative.**
- Purified antibodies: avoid repeated freeze/thaw cycles.**
- Coupled antibodies: glycerol protects from freezing.**

**Not for use in Humans. For research purpose only**